

Zebrafish as a Novel Vertebrate Model To Dissect Enterococcal Pathogenesis

Tomasz K. Prajsnar,^{a,b,c} Stephen A. Renshaw,^{a,c,d} Nikolay V. Ogryzko,^{a,c,d} Simon J. Foster,^{a,b} Pascale Serror,^{e,f} Stéphane Mesnage^{a,b}

Krebs Institute, University of Sheffield, Sheffield, United Kingdom^a; Department of Molecular Biology and Biotechnology, University of Sheffield, Sheffield, United Kingdom^b; MRC Centre for Developmental and Biomedical Genetics, University of Sheffield, Sheffield, United Kingdom^c; Department of Infection and Immunity, University of Sheffield, Sheffield, United Kingdom^d; INRA, UMR1319 Micalis, Jouy-en-Josas, France^e; AgroParisTech, UMR Micalis, Jouy-en-Josas, France^f

***Enterococcus faecalis* is an opportunistic pathogen responsible for a wide range of life-threatening nosocomial infections, such as septicemia, peritonitis, and endocarditis. *E. faecalis* infections are associated with a high mortality and substantial health care costs and cause therapeutic problems due to the intrinsic resistance of this bacterium to antibiotics. Several factors contributing to *E. faecalis* virulence have been identified. Due to the variety of infections caused by this organism, numerous animal models have been used to mimic *E. faecalis* infections, but none of them is considered ideal for monitoring pathogenesis. Here, we studied for the first time *E. faecalis* pathogenesis in zebrafish larvae. Using model strains, chosen isogenic mutants, and fluorescent derivatives expressing green fluorescent protein (GFP), we analyzed both lethality and bacterial dissemination in infected larvae. Genetically engineered immunocompromised zebrafish allowed the identification of two critical steps for successful establishment of disease: (i) host phagocytosis evasion mediated by the Epa rhamnopolysaccharide and (ii) tissue damage mediated by the quorum-sensing Fsr regulon. Our results reveal that the zebrafish is a novel, powerful model for studying *E. faecalis* pathogenesis, enabling us to dissect the mechanism of enterococcal virulence.**

E*nterococcus faecalis* is a Gram-positive commensal bacterium that colonizes the gastrointestinal tracts of humans and various animals. This bacterium is also responsible for a wide range of community- and hospital-acquired infections, including life-threatening bacteremia, infective endocarditis and peritonitis, and wound and urinary tract infections (1). The main risk factors for developing *E. faecalis* infections include impairment of the immune system, severe underlying diseases, urinary or vascular catheters, prior antibiotic therapy, and a prolonged stay in a hospital or intensive care unit (2). In some cases, diseases caused by *E. faecalis* are very difficult to treat, due to the large repertoire of intrinsic and acquired antibiotic resistance of this organism (3). Enterococci also represent a reservoir of genes conferring antibiotic resistance which can be disseminated to other pathogens, such as *Staphylococcus aureus* (4). Understanding the mechanisms of *E. faecalis* pathogenesis is therefore important to design preventive and alternative therapeutic approaches to treat enterococcal infections.

Several *E. faecalis* virulence factors have been identified using both mammalian [mainly mouse, rat, and rabbit (reviewed in reference 5)] and invertebrate [*Caenorhabditis elegans* (6), *Galleria mellonella* (7), and *Drosophila melanogaster* (8)] models of infection. These virulence factors are involved in attachment to host cells and extracellular matrix proteins, in cell and tissue damage and in immune system evasion (reviewed in references 3 and 9). Although invertebrates are amenable to large-scale genetic screens (10, 11), their physiology and immune systems differ significantly from mammalian systems. The zebrafish is a vertebrate organism that has been extensively used for developmental studies. The immune system of zebrafish shares with mammalian systems both innate and adaptive cellular immune system components (12), including macrophages and neutrophils already present at 1 day postfertilization (13). On a molecular level, high homology between zebrafish and human immune systems has also been observed, with Toll-like receptors (14, 15), nucleotide oligomeriza-

tion domain receptors NOD1 and NOD2 (16), and the complement system (17, 18). In addition, since zebrafish larvae are optically transparent, most cell types, including macrophages (19) and neutrophils (20), can be microscopically imaged in real time upon inoculation of fluorescent bacteria into the host. As a genetically tractable organism that can be obtained in high numbers, the zebrafish combines several advantages of both invertebrate and rodent infection models. It has been used to study virulence in *Salmonella enterica* (21), *Pseudomonas aeruginosa* (22, 23), *Burkholderia cenocepacia* (24), *S. aureus* (25, 26), *Listeria monocytogenes* (27), and *Streptococcus pneumoniae* (28), among others. In addition, studies on a natural fish pathogen—*Mycobacterium marinum*—using embryonic and adult zebrafish have provided multiple new insights into granuloma formation and disease progression of tuberculosis (29). The zebrafish has thus become a model of choice for studying bacterial human diseases.

In this study, we explore the relevance of a zebrafish model of infection to study *E. faecalis* virulence. Using a combinatorial approach, targeting both host and pathogen components to dissect the complex host-pathogen interactions, we found that the ze-

Received 8 August 2013 Returned for modification 25 August 2013

Accepted 29 August 2013

Published ahead of print 3 September 2013

Editor: A. J. Bäuml

Address correspondence to Stéphane Mesnage, s.mesnage@sheffield.ac.uk, or Tomasz K. Prajsnar, t.k.prajsnar@gmail.com.

Supplemental material for this article may be found at <http://dx.doi.org/10.1128/IAI.00976-13>.

Copyright © 2013 Prajsnar et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

doi:10.1128/IAI.00976-13

brafish model of infection is particularly suitable for studying both *E. faecalis*-induced lethality and pathogenesis.

MATERIALS AND METHODS

Bacterial strains, plasmids, and growth conditions. *E. faecalis* strains and plasmids are described in Table S1 in the supplemental material. Bacteria were grown in brain heart infusion (BHI) broth medium (Oxoid) at 37°C supplemented with antibiotics where appropriate at the following concentrations: tetracycline, 5 µg/ml; kanamycin, 2,000 µg/ml; and erythromycin, 30 µg/ml. *E. faecalis* strains were transformed with pMV158GFP (30) and pTEX5249 (31) by electroporation as previously described (31).

Zebrafish maintenance and transgenesis. London wild-type (LWT) inbred zebrafish embryos were provided by the aquarium staff of the MRC Centre for Developmental and Biomedical Genetics for zebrafish husbandry. The Tg(*mpeg1:Gal4.VP-16*)^{sh256} strain was generated as previously described (19). The Tg(*mpeg1:Gal4.VP-16*)^{sh256} strain was crossed to the Tg(*UAS:Kaede*)^{s1999t} strain to enable visualization of macrophages. Embryos were incubated in E3 medium at 28.5°C according to standard protocols (32).

Morpholino knockdown of *pu.1*. Morpholino-modified antisense oligomers against *pu.1* were injected as previously described (25).

Microinjections of *E. faecalis* into zebrafish embryos. Bacteria were grown in BHI broth until they reached an optical density at 600 nm of about 0.7 and harvested by centrifugation (5,500 × g, 10 min). Bacteria were microinjected into the circulation of dechorionated zebrafish embryos at 30 h postfertilization (hpf) as previously described (25). Briefly, anesthetized embryos were embedded in 3% (wt/vol) methylcellulose and injected individually using microcapillary pipettes filled with the bacterial suspension of known concentration. Larvae were observed frequently up to 90 h postinfection (hpi).

Microscopic observations of larvae. Live anesthetized larvae were immersed in 1% (wt/vol) low-melting-point agarose solution in E3 medium and mounted flat on a transparent slide. Images were acquired using the TE-2000U microscope (Nikon) with a Hamamatsu Orca-AG camera. Image acquisition and processing were performed with Volocity software (Improvision). A 4× Nikon Plan Fluor objective with a numerical aperture (NA) of 0.13 and a 60× Nikon Plan Apo oil objective with an NA of 1.4 were used.

Determination of *in vivo* bacterial load. At various times postinfection, six living zebrafish larvae were anesthetized and individually transferred with 100 µl of E3 medium into 0.5-ml Precellys tubes containing 1.4-mm ceramic beads (Peqlab) and homogenized using a Precellys 24-Dual homogenizer (Peqlab). The homogenates were serially diluted and plated on BHI agar to determine *E. faecalis* CFU numbers. Bacterial load was also determined for dead larvae at each time point.

Phagocytosis assay. Prior to infection, bacteria were labeled with pHrodo red succinimidyl ester (Invitrogen) by mixing 200 µl of bacterial suspension with 0.5 µl pHrodo red (2.5 mM) and incubating for 30 min in the dark. To remove the excess dye, bacteria were then washed with phosphate-buffered saline (PBS) followed by 50 mM Tris-HCl (pH 8.5) and subsequently resuspended in PBS. At 1.5 h postinfection (hpi), 18 larvae infected with each strain tested were mounted in 1% low-melting-point agarose, and microscopic images were captured using a 2× Nikon Plan UW objective (NA, 0.06) with a 543-nm excitation channel. The signal from pHrodo was then quantified for each embryo using ImageJ.

Analysis of exoproteins. *E. faecalis* strains were grown in BHI broth until exponential phase (optical density at 600 nm = 0.7) and harvested by centrifugation. Proteins in supernatants were precipitated by addition of trichloroacetic acid (TCA) at a 10% (wt/vol) final concentration. After incubation for 15 min on ice, proteins were recovered by centrifugation, and the protein pellet was washed with acetone. The final protein pellet was resuspended in Laemmli buffer prior to loading on a 12% (wt/vol) separation gel. The gel was stained with Coomassie blue R-250.




FIG 1 Intravenous infection of zebrafish embryos with *E. faecalis* OG1RF and V583 induces lethality, whereas infection with *E. faecalis* JH2-2 strain does not. (A) Survival of zebrafish larvae following injection with different doses of strains JH2-2, OG1RF, and V583 ($n = 25$). (B) Survival of zebrafish larvae following injection of various doses of *E. faecalis* OG1RF ($n = 25$). (C and D) Growth of *E. faecalis* OG1RF (C) and JH2-2 (D) within zebrafish larvae. (E) *In vivo* images of zebrafish larvae at 18 hpi following injection with GFP-labeled OG1RF. Bars, 500 µm. The numbers of CFU injected are indicated on each panel.

Statistical analyses. Survival experiments were evaluated using the Kaplan-Meier method. Comparisons between curves were made using the log rank test. The total pixel number in the phagocytosis assay was compared between two groups (OG1RF and *epaB*) using a two-tailed, unpaired Student's *t* test. Analysis was performed using Prism version 5.0 (GraphPad). Statistical significance was assumed at *P* values below 0.05.

RESULTS

Virulence of *E. faecalis* isolates in a zebrafish embryo infection model. We analyzed the virulence of three *E. faecalis* reference strains in a zebrafish embryo infection model: OG1RF and V583, two clinical isolates extensively used to study *E. faecalis* pathogenesis, and JH2-2, a nonvirulent laboratory strain used for genetic studies. Each strain was injected into the bloodstream of LWT zebrafish embryos at 30 h postfertilization (hpf). Embryos were maintained at 28.5°C, and the survival of larvae was monitored for 90 h postinfection (hpi). Injection of high numbers of CFU (10,500 to 12,000) showed that the three strains displayed radically distinct virulence phenotypes (Fig. 1A and B): while JH2-2 killed only 10% of larvae at 90 hpi, OG1RF killed 100% of larvae by 24 hpi. V583 displayed an intermediate virulence, killing 50% of larvae at 42 hpi. Injection of different numbers of CFU revealed that killing by *E. faecalis* was dose dependent. OG1RF was more virulent than V583, as 4,000 CFU of this strain were able to kill approximately 80% of larvae at 24 hpi, whereas 4,900 CFU of V583 killed only 10% of larvae. Remarkably, OG1RF was still vir-

ulent at relatively low doses (1,100 CFU induced 50% mortality at 42 hpi).

The striking differences between OG1RF and JH2-2 prompted us to enumerate bacteria within larvae following infection. Embryos infected with sublethal doses of either OG1RF or JH2-2 strain were homogenized, and the numbers of bacteria within individual larvae were counted. In larvae infected with OG1RF, the number of bacteria increased slightly during the first 7 hpi. In some hosts, numbers did not increase, and larvae survived. In several larvae, CFU increased exponentially, reaching approximately 10^6 CFU at 20 to 30 hpi, at which point death often occurred (Fig. 1C). The variable number of CFU from larva to larva may reflect the variability between individual hosts. Alternatively, we can speculate that high CFU numbers reflect a more advanced infection, leading to death when CFU numbers reach 10^5 to 10^6 per larva. In contrast, strain JH2-2 was unable to proliferate *in vivo*, and the bacterial numbers remained constant until at least 55 hpi (Fig. 1D).

Taking advantage of the embryonic optical transparency of developing zebrafish, we monitored bacterial dissemination during infection. For comparison purposes, both OG1RF and JH2-2 were transformed with a replicative plasmid constitutively expressing *gfp*. Zebrafish larvae were infected with one of the two *E. faecalis*-transformed strains and analyzed by fluorescence microscopy. Larvae infected with OG1RF developed lesions resembling necrotic tissues at sites of bacterial growth, referred to here as “tissue damage” (Fig. 1E, arrows). In contrast, the JH2-2-infected larvae remained apparently normal, with bacteria being restricted to small foci within the embryonic vasculature (data not shown). The OG1RF strain was able to disseminate systemically and proliferate within larvae and was often detected in the heart and central nervous system prior to embryo death (see Fig. S1A in the supplemental material). Zebrafish larvae that survived the infection were not able to completely clear bacteria from their bodies, and small foci of infection were seen until at least 90 hpi (see Fig. S1B).

Contribution of phagocytes to *E. faecalis* infection. At 30 hpf, embryo immune defenses rely on the innate response, primarily macrophages and neutrophils. To gain insight into the differences in virulence of OG1RF and JH2-2, we compared bacterial uptake by phagocytes at 2 hpi. In zebrafish larvae infected with 1,500 CFU of fluorescent *E. faecalis* OG1RF, a large proportion of fluorescent bacteria resisted phagocytosis and remained free in the bloodstream (Fig. 2A, arrows). In contrast, JH2-2 bacteria injected at a much higher dose (10,000 CFU) were all promptly phagocytosed (Fig. 2B).

To determine specific macrophage roles in engulfing injected enterococci, we used embryos expressing the macrophage-specific transgenes *mpeg1:Gal4* and *UAS:Kaede* (19). At 29 hpf, embryos were subjected to 436 nm irradiation for 20 min to enable Kaede photoconversion from green to red. The embryos were then infected with *gfp*-expressing *E. faecalis* JH2-2. Similarly to previous studies with other bacteria (25, 33), the majority of fluorescent enterococci were detected within macrophages (Fig. 2C).

To further explore the role of zebrafish phagocytes in resistance to *E. faecalis* infection, phagocytes were depleted using antisense *pu.1* morpholinos. *pu.1* morphants were significantly more susceptible to both OG1RF and JH2-2 than their immunocompetent counterparts (Fig. 2D). However, while OG1RF rapidly killed all phagocyte-depleted larvae, JH2-2 required 3 to 4 days to kill im-

munocompromised larvae. Despite the distinct death kinetics associated with strains OG1RF and JH2-2, both were able to proliferate rapidly in *pu.1* morphants, reaching approximately 10^6 CFU per embryo at 20 hpi (Fig. 2E and F). Imaging of *pu.1* morphants when bacterial proliferation was maximal showed serious deformation and tissue damage in OG1RF-infected larvae (Fig. 2G), whereas larvae infected with JH2-2 showed no signs of disease despite heavy dissemination (Fig. 2H). This result suggested that in addition to being easily phagocytosed, the JH2-2 strain lacks virulence determinants that contribute to killing of the host. Interestingly, this work indicates that zebrafish larvae represent a tractable model for studying *E. faecalis* pathogenesis under conditions that mimic neutropenia, a major risk factor for *E. faecalis* infections.

Contribution of *epa*, *fsr*, *gelE*, and *sprE* to *E. faecalis* pathogenesis. To gain further insights into the zebrafish model of *E. faecalis* infection, we analyzed OG1RF mutants previously shown to be attenuated in other animal models. Based on our results, which suggested a role of phagocytosis and revealed substantial tissue damage following infection, we focused on two types of virulence factors: (i) the enterococcal polysaccharide antigen (encoded by the *epa* locus), a rhamnopolysaccharide that has previously been shown to play a role in resistance to neutrophil uptake *in vitro* (34), and (ii) two major exoproteases playing a role in virulence, a gelatinase (*GelE*) and a serine protease (*SprE*) that are regulated by the *Fsr* quorum-sensing system (7, 31, 35). Four OG1RF mutants (*epaB*, *gelE*, *sprE*, and *fsr* mutants) were tested in our infection model.

The *epaB* mutant was significantly attenuated compared to the parental strain (Fig. 3A) (*P* value of 0.0001). To investigate the role of *Epa* in phagocytosis *in vivo*, we carried out microscopic examination of larvae infected with *epaB* mutant expressing green fluorescent protein (GFP). Unlike the parental strain (Fig. 3B), the *epaB* mutant was not able to evade phagocytosis *in vivo* and was readily taken up by zebrafish phagocytes compared to the wild-type strain (Fig. 3C). The difference in phagocytic uptake between OG1RF and its *epaB* derivative was quantified following labeling of bacterial cells with pHrodo-S-ester, a pH-sensitive dye that enables us to visualize bacteria only in low-pH compartments of phagosomes (36). This assay confirmed that at 1.5 hpi, the *epaB* mutant is phagocytosed more efficiently than the parental strain (Fig. 3D). Bacterial counts following infection with the *epaB* mutant remained constant until 55 hpi (Fig. 3E), clearly showing that although growth was abolished in zebrafish larvae, *epaB*-deficient bacteria were still able to survive within phagosomes. Together, our results therefore indicate that the *Epa* rhamnopolysaccharide contributes to evasion of phagocytosis but does not contribute to survival in zebrafish phagosomes.

In our model, *gelE*, *sprE*, and *fsr* mutants were significantly attenuated compared to the parental strain (Fig. 4A and B) (*P* values of 0.04, 0.03, and 0.0001 for *gelE*, *sprE*, and *fsr* mutants, respectively). Both *gelE* and *fsrB* mutants were selected to be further analyzed *in vivo*, as two examples of intermediate or severe reduction of virulence, respectively. Unexpectedly, both attenuated strains were able to proliferate similarly to the parental strain (Fig. 4C and D), although this growth did not result in larva death. In agreement with this observation, time-lapse imaging of larvae infected with *fsrB* mutants expressing *gfp* did not reveal any morphological change, even when the majority of the developing zebrafish were overwhelmed with bacteria (Fig. 4E). This obser-




FIG 2 Differential uptake of OG1RF and JH2-2 strains by phagocytes plays an important but not exclusive role in killing. (A and B) *In vivo* images of the yolk circulation valley of embryos at 2 hpi following injection with OG1RF (A) and JH2-2 (B) strains expressing *gfp*. Arrows indicate examples of unphagocytosed bacteria. These appear as “smears” due to their movement in the bloodstream and the relatively long exposure time. (C) *In vivo* imaging of the yolk circulation valley of Tg(*mpeg1*:Gal4 × UAS:*Kaede*) transgenic embryos at 2 hpi injected with GFP-expressing JH2-2. Photoconverted Kaede-positive macrophages (M) and nonfluorescent neutrophils (N) containing fluorescent *E. faecalis* are indicated. (D) Survival of wild-type or phagocyte-depleted zebrafish larvae following injection with *E. faecalis* OG1RF or JH2-2 ($n = 25$). (E and F) Growth of *E. faecalis* within phagocyte-depleted zebrafish larvae upon injection with OG1RF (E) or JH2-2 (F) strain into the bloodstream. (G and H) *In vivo* images of phagocyte-depleted zebrafish larvae injected with GFP-expressing OG1RF (G) or GFP-expressing JH2-2 at 18 hpi (H). Bars, 10 μ m (A, B, and C) and 500 μ m (G and H). The numbers of CFU injected are indicated on each panel.




FIG 3 Enterococcal polysaccharide antigen (*epa*) mutants of OG1RF are unable to evade phagocytosis and fail to establish lethal infection. (A) Survival of zebrafish larvae following injection with *E. faecalis* OG1RF and *epa* derivatives ($n = 25$). In log rank tests for *epaB* versus wild type, P is 0.0001. (B and C) *In vivo* images of the yolk circulation valley of larvae at 2 hpi following injection with *gfp*-expressing OG1RF (B) or the *epaB* mutant (C). Arrows indicate examples of unphagocytosed bacteria. (D) Efficacy of phagocytosis by zebrafish embryos ($n = 20$) at 1.5 h following infection with pHrodo-labeled wild-type *E. faecalis* OG1RF or the *epaB* mutant. (E) Growth of the *E. faecalis epaB* mutant in zebrafish larvae. The numbers of CFU injected are indicated on each panel. Bars, 10 μ m.

vation indicates that the Fsr system is not required for *in vivo* proliferation but contributes to late stages of infection.

To determine whether the attenuation of the mutants analyzed was caused by their inability to overcome the host cellular immune response, we followed the infection of phagocyte-depleted larvae (Fig. 5). As expected, the virulence of *epaB* mutant was fully restored in *pu.1* morphants. In contrast, death of larvae infected with the *gelE* mutants was significantly delayed compared to that of the parental strain. For *pu.1* morphants infected with the *fsrB* mutant, the time to death was severely delayed (Fig. 5A). Interestingly, bacterial growth in larvae was very similar for all mutants tested, reaching approximately 10^6 CFU at 20 hpi (Fig. 5B to D). However, different death rates indicated that *in vivo* proliferation is not sufficient to cause death. Altogether, these results indicate that zebrafish represent a particularly suitable model for following two steps crucial to the progression of *E. faecalis* pathogenesis: phagocytosis evasion and host killing, in part mediated by exoprotease genes and other *fsrB*-dependent genes.

Heterologous complementation of the *fsr* locus in JH2-2. Survival of *pu.1* knockdown larvae infected with the OG1RF *fsrB*

mutant (Fig. 5A) was very similar to that of immunocompromised larvae infected with JH2-2 (Fig. 2D). We hypothesized that the inactive *fsr* system in JH2-2 could at least in part account for the delay in mortality in immunocompromised larvae. To test this hypothesis, we complemented the JH2-2 strain with the OG1RF *fsr* locus using plasmid pTEX5249 (31). As anticipated, heterologous expression of the OG1RF *fsr* locus restored the production of exoproteases in JH2-2 (37) (see Fig. S2 in the supplemental material). JH2-2 *fsr*⁺ bacteria were avirulent when injected into immunocompetent larvae but rapidly killed phagocyte-depleted larvae (Fig. 6A). Analysis of bacterial load following infection revealed no growth of JH2-2 *fsr*⁺ bacteria in wild-type larvae (Fig. 6B), indicating that restoration of neither exoprotease production nor any other Fsr-dependent factor allowed JH2-2 to overcome phagocytosis. In *pu.1* knockdown larvae, JH2-2 *fsr*⁺ bacteria proliferated rapidly and caused death of infected larvae (Fig. 6C), highlighting the important role of Fsr-dependent factors, including exoproteases, in later stages of infection, after the initial immune hurdle (i.e., phagocytes) has been overcome. These results show that the genetically manipulable strain JH2-2 can be used as a platform strain to investigate functions of virulence factors.

DISCUSSION

Several vertebrate models for studying specific *E. faecalis* infections have been described: endophthalmitis and endocarditis in rabbits and urinary tract infection, bacteremia, and peritonitis in mice (5). Although these models have been successfully used to characterize a large variety of *E. faecalis* virulence factors, they are not adapted for large-scale studies and remain costly. Over the past decade, zebrafish have been successfully used as a surrogate model host for human bacterial pathogens (38). In this study, we showed that zebrafish can be used as a host to study *E. faecalis* pathogenesis. Our results revealed that this infection model is particularly suitable for identifying *E. faecalis* virulence factors, dissecting the mechanisms of infection, and following the dissemination of bacteria *in vivo*.

Zebrafish as a model for studying *E. faecalis* virulence. This work showed that zebrafish larvae are susceptible to virulent *E. faecalis* strains (OG1RF and V583) in a dose-dependent manner but resistant to strain JH2-2, a nonvirulent laboratory strain. In agreement with previous studies comparing OG1RF and V583 in mouse and *C. elegans* models (6, 39), we also show that OG1RF was more virulent than V583 in the zebrafish lethality model (Fig. 1). This work also confirms the poor pathogenic potential of JH2-2. When high numbers of CFU of these bacteria were injected, they failed to induce lethality despite the fact that they were able to multiply and persist *in vivo*.

To demonstrate the relevance of the zebrafish model of infection, we focused on the analysis of mutations in the protease genes *gelE* and *sprE* and the *fsr* quorum-sensing system, which positively controls their expression (31). Mutation in the *fsr* locus has been associated with a significant decrease in virulence in all animal models tested (7, 31, 35, 40). Similarly to previous studies on various animal models, we found that *fsrB* deletion resulted in severe loss of enterococcal virulence in zebrafish larvae. However, we showed that the *fsrB* mutant was able to proliferate as well as its parent strain, OG1RF (Fig. 4), indicating that *fsrB* controls factors important for the late stages of infection, leading to killing. While *GelE* has been un-




FIG 4 Protease-deficient mutants of OG1RF proliferate *in vivo* but are less virulent than the wild type. (A and B) Survival of zebrafish larvae ($n \geq 36$) following injection with the indicated *E. faecalis* mutant strains. In log rank tests, for the *gelE* mutant versus the wild type (WT), $P = 0.04$; for the *fsrB* mutant versus the WT, $P = 0.0001$; for the *sprE* mutant versus the WT, $P = 0.03$; for the *sprE gelE* double mutant versus the WT, $P = 0.0006$. (C and D) Growth of the *E. faecalis gelE* (C) or *fsrB* (D) mutant in zebrafish larvae (E). Time-lapse microscopy of a representative zebrafish larva infected with a GFP-expressing *E. faecalis fsrB* mutant. The numbers of CFU injected are indicated on each panel. Bars, 500 μm .

ambiguously demonstrated to be important for enterococcal virulence in multiple models (31, 35, 41–43), the importance of SprE as an *E. faecalis* virulence factor is controversial. The *sprE* mutant was shown to be attenuated in mouse peritonitis in the nematode and the fly models (8, 31, 35). However, SprE was dispensable for rabbit infective endocarditis and had no lethal activity in *Galleria mellonella* (42, 43). The role of these exoproteases in promoting tissue damage is strongly supported based on histopathological examination of infected eyes using the endophthalmitis model (44) and their proteolytic activity over a wide substrate range (45). Here, we showed that both the gelatinase GelE and the serine protease SprE contribute to *E. faecalis* virulence in zebrafish larvae and appear to have additive effects in pathogenesis. Altogether, given that *E. faecalis* is able to proliferate and disseminate within zebrafish larvae, this infection model appears to be suitable for identifying a range of virulence factors associated with *E. faecalis* pathogenesis. One potential limitation of the zebrafish infection model is the temperature at which the larvae are kept (28.5°C). Thus, alternative infection models need to be considered to analyze virulence

factors specifically expressed at 37°C, the temperature of the mammalian host.

Zebrafish as a model system to study innate immune response against *E. faecalis*. In this study, we showed that zebrafish larvae can be used as a model to evaluate both uptake and survival of bacteria in phagocytes. Upon injection into the bloodstream, *E. faecalis* phagocytosis is mediated by macrophages. This result is in agreement with previous studies indicating that in zebrafish, bacteria in blood circulation are essentially phagocytosed by macrophages, neutrophils being able to engulf only surface-associated microbes (46). Although macrophages completely phagocytosed a large inoculum of JH2-2, a significant proportion of OG1RF cells escaped phagocytosis and remained in the bloodstream. *E. faecalis*-specific rhamnopolysaccharide encoded by the *epa* locus was previously shown to mediate resistance to uptake and killing by human neutrophils (11, 34). Using the *Epa*-deficient *epaB* mutant (47), we showed that the *Epa* rhamnopolysaccharide protects OG1RF from uptake by macrophages in zebrafish. In the phagocyte-depleted larvae, the *epaB* mutant was as virulent as




FIG 5 Protease-deficient mutants of OG1RF remain attenuated in phagocyte-depleted larvae. (A) Survival of *pu.1* knockdown zebrafish larvae ($n = 25$) following infection with *E. faecalis* *fsrB*, *gelE*, and *epaB* mutants. In log rank tests, for *gelE* versus WT, $P < 0.0001$; for *fsrB* versus WT, $P < 0.0001$; for *epaB* versus WT, $P = 0.97$. (B to D) Growth of *E. faecalis* *epaB* (B), *gelE* (C), and *fsrB* (D) mutants in *pu.1* knockdown larvae.

the parental OG1RF strain, indicating that during infection, the main role of the *epa* cluster is to prevent phagocytosis. However, our data indicate that the rhamnopolysaccharide encoded by the JH2-2 *epa* locus is not able to confer resistance to phagocytosis *in vivo*. Diversity of the *epa* locus within *E. faecalis* species was recently reported (48). The zebrafish model is particularly appropriate for evaluating whether *Epa* diversity impacts *E. faecalis* resistance to phagocytosis *in vivo*. *E. faecalis* persistence within zebrafish larvae is in agreement with the extreme resistance of this bacterium to various stresses (reviewed in reference 49). It will be interesting to test whether factors that have been shown to enhance survival in murine macrophages (50) or human neutrophils (51) play a role in resistance to zebrafish macrophages. Neutropenia is a major risk factor for enterococcal infections (2). The higher susceptibility of immunocompromised zebrafish larvae to *E. faecalis*

infection suggests that this model is suitable for investigating *E. faecalis* pathogenesis under conditions mimicking disease onset.

Concluding remarks. Our study revealed that zebrafish larvae represent an attractive alternative model system for studying host-pathogen interaction during *E. faecalis* infections, allowing the identification of both host immune components and pathogen virulence factors. This novel vertebrate model enables spatial and temporal dissection of the complex mechanisms of *E. faecalis* infection *in vivo*, from initial interaction with host innate immune system to development of bacterial quorum sensing *in vivo* and degradation of host tissues, resulting in death. We showed that the degree of virulence can be readily measured by a simple lethality assay. More importantly, thanks to zebrafish optical transparency and genetic tractability, we also reveal that successful infection by *E. faecalis* relies




FIG 6 Heterologous expression of OG1RF *fsr* locus in JH2-2 restores virulence in phagocyte-depleted but not wild-type larvae. (A) Survival of wild-type or *pu.1* knockdown zebrafish larvae ($n = 25$) following infection with *E. faecalis* JH2-2 with or without *fsrB* complementation. In log rank tests, for JH2-2 versus JH2-2 *fsr*⁺ in wild-type larvae, $P = 0.56$; for JH2-2 versus JH2-2 *fsr*⁺ in *pu.1* knockdown larvae, $P < 0.0001$. (B and C) Growth of *E. faecalis* JH2-2 *fsr*⁺ in wild-type (B) or *pu.1* knockdown (C) larvae. The numbers of CFU injected are indicated on each panel.

on two major steps: phagocyte evasion and tissue damage. Based on these observations, we are confident that the model described here will be useful for large-scale screening of mutant libraries to identify new bacterial virulence determinants and host factors important in immunity against enterococci.

ACKNOWLEDGMENTS

We thank Cristel Archambaud for useful comments on the manuscript, Barbara E. Murray and Christopher J. Kristich for providing mutant strains used in this study, and Justyna Serba for sharing the bacterial staining protocol with pHrodo.

This work was funded by a Wellcome Trust Project Grant (reference number WT089981MA) and the European Marie-Curie Initial Training Network FishForPharma (PITN-GA-2011-289209).

REFERENCES

- Murray BE. 1990. The life and times of the *Enterococcus*. Clin. Microbiol. Rev. 3:46–65.
- de Perio MA, Yarnold PR, Warren J, Noskin GA. 2006. Risk factors and outcomes associated with non-*Enterococcus faecalis*, non-*Enterococcus faecium* enterococcal bacteremia. Infect. Control Hosp. Epidemiol. 27: 28–33.
- Arias CA, Murray BE. 2012. The rise of the *Enterococcus*: beyond vancomycin resistance. Nat. Rev. Microbiol. 10:266–278.
- Weigel LM, Clewell DB, Gill SR, Clark NC, McDougal LK, Flannagan SE, Kolonay JF, Shetty J, Killgore GE, Tenover FC. 2003. Genetic analysis of a high-level vancomycin-resistant isolate of *Staphylococcus aureus*. Science 302:1569–1571.
- Tendolkar PM, Baghdayan AS, Shankar N. 2003. Pathogenic enterococci: new developments in the 21st century. Cell Mol. Life Sci. 60:2622–2636.
- Garsin DA, Sifri CD, Mylonakis E, Qin X, Singh KV, Murray BE, Calderwood SB, Ausubel FM. 2001. A simple model host for identifying Gram-positive virulence factors. Proc. Natl. Acad. Sci. U. S. A. 98:10892–10897.
- Gaspar F, Teixeira N, Rigottier-Gois L, Marujo P, Nielsen-LeRoux C, Crespo MTB, de Fátima Silva Lopes M, Serror P. 2009. Virulence of *Enterococcus faecalis* dairy strains in an insect model: the role of *fsrB* and *gelE*. Microbiology 155:3564–3571.
- Teixeira N, Varahan S, Gorman MJ, Palmer KL, Zaidman-Remy A, Yokohata R, Nakayama J, Hancock LE, Jacinto A, Gilmore MS, de Fátima Silva Lopes M. 2013. Drosophila host model reveals new *Enterococcus faecalis* quorum-sensing associated virulence factors. PLoS One 8:e64740. doi:10.1371/journal.pone.0064740.
- Sava IG, Heikens E, Huebner J. 2010. Pathogenesis and immunity in enterococcal infections. Clin. Microbiol. Infect. 16:533–540.
- Garsin DA, Urbach J, Huguet-Tapia JC, Peters JE, Ausubel FM. 2004. Construction of an *Enterococcus faecalis* *Tn917*-mediated-gene-disruption library offers insight into *Tn917* insertion patterns. J. Bacteriol. 186:7280–7289.
- Rigottier-Gois L, Alberti A, Houel A, Taly J-F, Palcy P, Manson J, Pinto D, Matos RC, Carrilero L, Montero N, Tariq M, Karsens H, Repp C, Kropec A, Budin-Verneuil A, Benachour A, Sauvageot N, Bizzini A, Gilmore MS, Bessières P, Kok J, Huebner J, Lopes F, Gonzalez-Zorn B, Hartke A, Serror P. 2011. Large-scale screening of a targeted *Enterococcus faecalis* mutant library identifies envelope fitness factors. PLoS One 6:e29023. doi:10.1371/journal.pone.0029023.
- Yoder JA, Nielsen ME, Amemiya CT, Litman GW. 2002. Zebrafish as an immunological model system. Microbes Infect. 4:1469–1478.
- Henry KM, Loynes CA, Whyte MKB, Renshaw SA. 2013. Zebrafish as a model for the study of neutrophil biology. J. Leukoc. Biol. [Epub ahead of print.] doi:10.1189/jlb.1112594.
- Meijer AH, Krens SFG, Medina Rodriguez IA, He S, Bitter W, Snaar-Jagalska BE, Spaik HP. 2004. Expression analysis of the Toll-like receptor and TIR domain adaptor families of zebrafish. Mol. Immunol. 40:773–783.
- van der Sar AM, Stockhammer OW, van der Laan C, Spaik HP, Bitter W, Meijer AH. 2006. MyD88 innate immune function in a zebrafish embryo infection model. Infect. Immun. 74:2436–2441.
- Oehlers SH, Flores MV, Hall CJ, Swift S, Crosier PS. 2011. The inflammatory bowel disease (IBD) susceptibility genes *NOD1* and *NOD2* have conserved anti-bacterial roles in zebrafish. Dis. Model. Mech. 4:832–841.
- Wang Z, Zhang S, Tong Z, Li L, Wang G. 2009. Maternal transfer and protective role of the alternative complement components in zebrafish *Danio rerio*. PLoS One 4:e4498. doi:10.1371/journal.pone.0004498.
- Wang Z, Zhang S, Wang G, An Y. 2008. Complement activity in the egg cytosol of zebrafish *Danio rerio*: evidence for the defense role of maternal complement components. PLoS One 3:e1463. doi:10.1371/journal.pone.0001463.
- Ellett F, Pase L, Hayman JW, Andrianopoulos A, Lieschke GJ. 2011. *mpeg1* promoter transgenes direct macrophage-lineage expression in zebrafish. Blood 117:e49–e56.
- Renshaw SA, Loynes CA, Trushell DM, Elworthy S, Ingham PW, Whyte MK. 2006. A transgenic zebrafish model of neutrophilic inflammation. Blood 108:3976–3978.
- van der Sar AM, Musters RJ, van Eeden FJ, Appelmelk BJ, Vandenbroucke-Grauls CM, Bitter W. 2003. Zebrafish embryos as a model host for the real time analysis of *Salmonella typhimurium* infections. Cell Microbiol. 5:601–611.
- Brannon MK, Davis JM, Mathias JR, Hall CJ, Emerson JC, Crosier PS, Huttenlocher A, Ramakrishnan L, Moskowitz SM. 2009. *Pseudomonas aeruginosa* type III secretion system interacts with phagocytes to modulate systemic infection of zebrafish embryos. Cell Microbiol. 11:755–768.
- Clatworthy AE, Lee JS, Leibman M, Kostun Z, Davidson AJ, Hung DT. 2009. *Pseudomonas aeruginosa* infection of zebrafish involves both host and pathogen determinants. Infect. Immun. 77:1293–1303.
- Vergunst AC, Meijer AH, Renshaw SA, O'Callaghan D. 2010. *Burkholderia cenocepacia* creates an intramacrophage replication niche in zebrafish embryos, followed by bacterial dissemination and establishment of systemic infection. Infect. Immun. 78:1495–1508.
- Prajnsar TK, Cunliffe VT, Foster SJ, Renshaw SA. 2008. A novel vertebrate model of *Staphylococcus aureus* infection reveals phagocyte-dependent resistance of zebrafish to non-host specialized pathogens. Cell Microbiol. 10:2312–2325.
- Prajnsar TK, Hamilton R, Garcia-Lara J, McVicker G, Williams A, Boots M, Foster SJ, Renshaw SA. 2012. A privileged intraphagocyte niche is responsible for disseminated infection of *Staphylococcus aureus* in a zebrafish model. Cell Microbiol. 14:1600–1619.
- Levrud J-P, Disson O, Kissa K, Bonne I, Cossart P, Herbomel P, Lecuit M. 2009. Real-time observation of *Listeria monocytogenes*-phagocyte interactions in living zebrafish larvae. Infect. Immun. 77:3651–3660.
- Rounioja S, Saralahti A, Rantala L, Parikka M, Henriques-Normark B, Silvennoinen O, Rämetsä M. 2012. Defense of zebrafish embryos against *Streptococcus pneumoniae* infection is dependent on the phagocytic activity of leukocytes. Dev. Comp. Immunol. 36:342–348.
- Berg RD, Ramakrishnan L. 2012. Insights into tuberculosis from the zebrafish model. Trends Mol. Med. 18:689–690.
- Nieto C, Espinosa M. 2003. Construction of the mobilizable plasmid pMV158GFP, a derivative of pMV158 that carries the gene encoding the green fluorescent protein. Plasmid 49:281–285.
- Qin X, Singh KV, Weinstock GM, Murray BE. 2000. Effects of *Enterococcus faecalis* *fsr* genes on production of gelatinase and a serine protease and virulence. Infect. Immun. 68:2579–2586.
- Nusslein-Volhard CDR. 2002. Zebrafish. A practical approach. Oxford University Press, New York, NY.
- Le Guyader D, Redd MJ, Colucci-Guyon E, Murayama E, Kissa K, Briolat V, Mordet E, Zapata A, Shinomiya H, Herbomel P. 2008. Origins and unconventional behavior of neutrophils in developing zebrafish. Blood 111:132–141.
- Teng F, Jacques-Palaz KD, Weinstock GM, Murray BE. 2002. Evidence that the enterococcal polysaccharide antigen gene (*epa*) cluster is widespread in *Enterococcus faecalis* and influences resistance to phagocytic killing of *E. faecalis*. Infect. Immun. 70:2010–2015.
- Sifri CD, Mylonakis E, Singh KV, Qin X, Garsin DA, Murray BE, Ausubel FM, Calderwood SB. 2002. Virulence effect of *Enterococcus faecalis* protease genes and the quorum sensing locus *fsr* in *Caenorhabditis elegans* and mice. Infect. Immun. 70:5647–5650.
- Miksa M, Komura H, Wu R, Shah KG, Wang P. 2009. A novel method to determine the engulfment of apoptotic cells by macrophages using pHrodo succinimidyl ester. J. Immunol. Methods 342:71–77.
- Shankar J, Walker RG, Ward D, Horsburgh MJ. 2012. The *Enterococcus*

- faecalis* exoproteome: identification and temporal regulation by Fsr. *PLoS One* 7:e33450. doi:10.1371/journal.pone.0033450.
38. Meijer AH, Spaink HP. 2011. Host-pathogen interactions made transparent with the zebrafish model. *Curr. Drug Targets* 12:1000–1017.
 39. Bourgogne A, Garsin DA, Qin X, Singh KV, Sillanpaa J, Yerrapragada S, Ding Y, Dugan-Rocha S, Buhay C, Shen H, Chen G, Williams G, Muzny D, Maadani A, Fox KA, Gioia J, Chen L, Shang Y, Arias CA, Nallapareddy SR, Zhao M, Prakash VP, Chowdhury S, Jiang H, Gibbs RA, Murray BE, Highlander SK, Weinstock GM. 2008. Large scale variation in *Enterococcus faecalis* illustrated by the genome analysis of strain OG1RF. *Genome Biol.* 9:R110.
 40. Mylonakis E, Engelbert M, Qin X, Sifri CD, Murray BE, Ausubel FM, Gilmore MS, Calderwood SB. 2002. The *Enterococcus faecalis* *fsrB* gene, a key component of the *fsr* quorum-sensing system, is associated with virulence in the rabbit endophthalmitis model. *Infect. Immun.* 70:4678–4681.
 41. Singh KV, Qin X, Weinstock GM, Murray BE. 1998. Generation and testing of mutants of *Enterococcus faecalis* in a mouse peritonitis model. *J. Infect. Dis.* 178:1416–1420.
 42. Park SY, Kim KM, Lee JH, Seo SJ, Lee IH. 2007. Extracellular gelatinase of *Enterococcus faecalis* destroys a defense system in insect hemolymph and human serum. *Infect. Immun.* 75:1861–1869.
 43. Thurlow LR, Thomas VC, Narayanan S, Olson S, Fleming SD, Hancock LE. 2010. Gelatinase contributes to the pathogenesis of endocarditis caused by *Enterococcus faecalis*. *Infect. Immun.* 78:4936–4943.
 44. Engelbert M, Mylonakis E, Ausubel FM, Calderwood SB, Gilmore MS. 2004. Contribution of gelatinase, serine protease, and *fsr* to the pathogenesis of *Enterococcus faecalis* endophthalmitis. *Infect. Immun.* 72:3628–3633.
 45. Mäkinen PL, Clewell DB, An F, Mäkinen KK. 1989. Purification and substrate specificity of a strongly hydrophobic extracellular metalloendopeptidase (“gelatinase”) from *Streptococcus faecalis* (strain OG1-10). *J. Biol. Chem.* 264:3325–3334.
 46. Colucci-Guyon E, Tinevez J-Y, Renshaw SA, Herbomel P. 2011. Strategies of professional phagocytes *in vivo*: unlike macrophages, neutrophils engulf only surface-associated microbes. *J. Cell Sci.* 124:3053–3059.
 47. Xu Y, Singh KV, Qin X, Murray BE, Weinstock GM. 2000. Analysis of a gene cluster of *Enterococcus faecalis* involved in polysaccharide biosynthesis. *Infect. Immun.* 68:815–823.
 48. Palmer KL, Godfrey P, Griggs A, Kos VN, Zucker J, Desjardins C, Cerqueira G, Gevers D, Walker S, Wortman J, Feldgarden M, Haas B, Birren B, Gilmore MS. 2012. Comparative genomics of enterococci: variation in *Enterococcus faecalis*, clade structure in *Enterococcus faecium*, and defining characteristics of *Enterococcus gallinarum* and *Enterococcus casseliflavus*. *mBio* 3:e00318–11. doi:10.1128/mBio.00318-11.
 49. Fisher K, Phillips C. 2009. The ecology, epidemiology and virulence of *Enterococcus*. *Microbiology* 155:1749–1757.
 50. Hébert L, Courtin P, Torelli R, Sanguinetti M, Chapot-Chartier M-P, Auffray Y, Benachour A. 2007. *Enterococcus faecalis* constitutes an unusual bacterial model in lysozyme resistance. *Infect. Immun.* 75:5390–5398.
 51. Rakita RM, Vanek NN, Jacques-Palaz K, Mee M, Mariscalco MM, Dunny GM, Snuggs M, Van Winkle WB, Simon SI. 1999. *Enterococcus faecalis* bearing aggregation substance is resistant to killing by human neutrophils despite phagocytosis and neutrophil activation. *Infect. Immun.* 67:6067–6075.